

Some Notes Related to Module EN0311 (Control Part)
From *Manuel Kuehner* (German)
2006/2007

Contents

1	Matrix algebra	4
1.1	Basic definitions	4
1.1.1	Order	4
1.1.2	Elements	4
1.2	Addition, subtraction scalar- and matrix multiplication	4
1.2.1	Addition and subtraction	4
1.2.2	Scalar multiplications	5
1.2.3	Matrix multiplications	5
1.3	The transpose of a matrix	6
1.3.1	Definition and properties	6
1.3.2	Example(s)	6
1.4	Special square matrices	7
1.4.1	Square matrices	7
1.4.2	Diagonal matrices	7
1.4.3	Identity matrices	7
1.4.4	Symmetric matrices	8
1.5	Determinants, Minor and Cofactor	8
1.5.1	Definition and properties	8
1.5.2	Example(s)	9
1.6	Adjoint	11
1.6.1	Definition and properties	11
1.6.2	Example(s)	11
1.7	Inverse matrices	13
1.7.1	Definition and properties	13
1.8	Eigenvalue	13
1.8.1	Definition and properties	13
1.8.2	Example(s)	14

2	Linear equations	14
3	Laplace transformation	15
3.1	Definition	15
3.2	A few important properties	15
3.3	Table of important transforms	17
4	State space	18
4.1	How to transform a n th order differential equation into a set of first order differential equations	18
4.2	Solution of a linear system	20
4.3	State-space to transfer function	20
5	IO to transfer function	21

1 Matrix algebra

1.1 Basic definitions

1.1.1 Order

Definition

An $m \times n$ matrix has m rows and n columns. The matrix has **order** $r \times c$ (german: Höhe mal Breite).

Example

$$A = \begin{pmatrix} 1 & 8 \\ 7 & 2 \\ 3 & 4 \end{pmatrix}$$

Matrix A has **order** 3×2 .

1.1.2 Elements

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}$$

Matrix A has order $m \times n$. a_{ij} represents the **element** in the i th row and j th column.

1.2 Addition, subtraction scalar- and matrix multiplication

1.2.1 Addition and subtraction

Definitions and properties

- Two matrices can be **added or subtracted** if they have the **same order**.
- Their **sum or difference** is found by **adding or subtracting corresponding elements**.
- Matrix addition is **commutative**: $A + B = B + A$
- Matrix addition is **associative**: $A + (B + C) = (A + B) + C$
- Matrix addition is **distributive** (see 1.2.2): $k(A + B) = kA + kB$, k scalar

Example(s)

$$A = \begin{pmatrix} 1 & 5 & -2 \\ 3 & 1 & 1 \end{pmatrix} \text{ and } B = \begin{pmatrix} 1 & 2 & 0 \\ 1 & 1 & 4 \end{pmatrix}$$

$$A + B = \begin{pmatrix} 1+1 & 5+2 & -2+0 \\ 3+1 & 1+1 & 1+4 \end{pmatrix} = \begin{pmatrix} 2 & 7 & -2 \\ 4 & 2 & 5 \end{pmatrix} = B + A$$

1.2.2 Scalar multiplications

$$\text{if } A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix} \text{ then } kA = \begin{pmatrix} ka_{11} & ka_{12} & \cdots & ka_{1n} \\ ka_{21} & ka_{22} & \cdots & ka_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ ka_{m1} & ka_{m2} & \cdots & ka_{mn} \end{pmatrix}, k \text{ scalar}$$

1.2.3 Matrix multiplications**Definition and properties**

- If A is $m \times n$ and B is $r \times s$ the **product** of AB can **only** be formed if $n = r$. The **product is** then a $m \times s$ matrix.
- If A is a $m \times p$ matrix and B is a $p \times n$ matrix then the product $C = AB$ is defined as the $m \times n$ matrix with elements:

$$c_{ij} = \sum_{k=1}^p a_{ik}b_{kj}, \quad i = 1, 2, \dots, m; \quad j = 1, 2, \dots, n$$

- Or you can say: To find c_{ij} you take the i th row of A and pair its elements with the j th column of B . The paired elements are multiplied together and added to form c_{ij} .
- $C = AB \rightarrow B$ has been **premultiplied** by A , or alternatively A has been **postmultiplied** by B .
- In general $AB \neq BA$ and so matrix multiplication is **not commutative**.
- Matrix multiplication is **associative** and so: $(AB)C = A(BC)$.

Example(s)

$$A = \begin{pmatrix} a & b \\ c & d \\ e & f \end{pmatrix} = (3 \times 2) \text{ and } B = \begin{pmatrix} g & h & i & j \\ k & l & m & n \end{pmatrix} = (2 \times 4)$$

Reminder: To find c_{ij} you take the i th row of A and pair its elements with the j th column of B . The paired elements are multiplied together and added to form c_{ij} .

$$C = \begin{pmatrix} c_{11} = ag + bk & c_{12} = ah + bl & c_{13} = ai + bm & c_{14} = aj + bn \\ c_{21} = cg + dk & c_{22} = ch + dl & c_{23} = ci + dm & c_{24} = cj + dn \\ c_{31} = eg + fk & c_{32} = eh + fl & c_{33} = ei + fm & c_{34} = ej + fn \end{pmatrix} = 3 \times 4$$

What happens when there are more than two multiplicands?

$$X = ABC$$

1. Calculate the product $U = BC$.
2. Calculate the product $X = AU$.

1.3 The transpose of a matrix

1.3.1 Definition and properties

If A is an arbitrary $m \times n$ matrix, a related matrix is the **transpose** of A , written A^T . The transpose A^T is found by **interchanging the rows and columns** of A . A^T is an $n \times m$ matrix.

- $(A + B)^T = A^T + B^T$
- $(kA)^T = kA^T$, k scalar
- $(A^T)^T = A$
- $(AB)^T = B^T A^T$ (Watch out!)
- $(A^{-1})^T = (A^T)^{-1}$

1.3.2 Example(s)

$$A = \begin{pmatrix} 4 & 2 & 6 \\ 1 & 8 & 7 \end{pmatrix} \rightarrow A^T = \begin{pmatrix} 4 & 1 \\ 2 & 8 \\ 6 & 7 \end{pmatrix}$$

1.4 Special square matrices

1.4.1 Square matrices

Definition

A matrix which has the **same number of rows as columns** is called **square** matrix.

Example(s)

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} = 3 \times 3 = m \times m$$

1.4.2 Diagonal matrices

Definition

A square matrix which has elements that are **zero everywhere except of the leading diagonal** is called a **diagonal** matrix.

Example(s)

Matrices A and B are diagonal matrices.

$$A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 8 \end{pmatrix} \quad B = \begin{pmatrix} 0 & 0 \\ 0 & 3 \end{pmatrix}$$

1.4.3 Identity matrices

Definition and property

Diagonal matrices which have **only ones on their leading diagonal** are called a **identity** matrices and are denoted by the letter I .

If A is a square matrix then $IA = A$

Example(s)

Matrices A and B are identity matrices.

$$A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} B = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

1.4.4 Symmetric matrices

Definition and property

If a square matrix A and its transpose A^T are identical, then A is said to be a **symmetric** matrix.

A symmetric matrix is symmetrical about its leading diagonal

Example(s)

$$A = \begin{pmatrix} 5 & -4 & 2 \\ -4 & 6 & 9 \\ 2 & 9 & 13 \end{pmatrix} = \begin{pmatrix} \dots & -4 & 2 \\ -4 & \dots & 9 \\ 2 & 9 & \dots \end{pmatrix} \rightarrow A^T = \begin{pmatrix} 5 & -4 & 2 \\ -4 & 6 & 9 \\ 2 & 9 & 13 \end{pmatrix} = A$$

1.5 Determinants, Minor and Cofactor

1.5.1 Definition and properties

The determinant of a **square** $n \times n$ matrix A is denoted by $\det(A)$ or $|A|$. If we take a determinant and delete row i and column j then the determinant remaining is called **minor** M_{ij} of the element a_{ij} . In general we can take any row i (or column j) and evaluate an $n \times n$ determinant $|A|$ as:

$$|A| = \sum_{j=1}^n \underbrace{(-1)^{i+j}}_{\text{sign}} a_{ij} \underbrace{M_{ij}}_{\text{minor}}$$

With **cofactor** $A_{ij} = (-1)^{i+j} M_{ij}$ (appropriate sign \times minor):

$$|A| = \sum_{j=1}^n a_{ij} A_{ij}$$

- $|A^T| = |A|$
- $|AB| = |A||B|$
- $|k| = k$, k scalar

- A square matrix A is said to be **non-singular** if $|A| \neq 0$ and **singular** if $|A| = 0$.
- If there is a linear combination (rows or columns) in the matrix then the determinant of the matrix is always zero.
- If the elements in the upper or lower triangle of the matrix are all zero then the determinant of the matrix is the product of the elements on the leading diagonal.

$(-1)^{i+j}$ can be depicted as:

$$\begin{pmatrix} + & - & + & \dots \\ - & + & - & \dots \\ + & - & + & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix} = \begin{pmatrix} (-1)^2 & (-1)^3 & (-1)^4 & \dots \\ (-1)^3 & (-1)^4 & (-1)^5 & \dots \\ (-1)^4 & (-1)^5 & (-1)^6 & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

The determinant of 2×2 matrix is calculated as follows:

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = \det \begin{pmatrix} a & b \\ c & d \end{pmatrix} = ad - bc$$

1.5.2 Example(s)

Expansion along the **first row** $\rightarrow i = 1$

$$A = \begin{pmatrix} 2 & -1 & 4 \\ 4 & 0 & 5 \\ -6 & 2 & 3 \end{pmatrix} = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$$

$$|A| = \det(A) = \begin{vmatrix} 2 & -1 & 4 \\ 4 & 0 & 5 \\ -6 & 2 & 3 \end{vmatrix}$$

With $i = 1$ and $n = 3$:

$$\begin{aligned} |A| &= \sum_{j=1}^3 \underbrace{(-1)^{1+j}}_{\text{sign}} a_{1j} \underbrace{M_{1j}}_{\text{minor}} \\ &= (-1)^2 a_{11} M_{11} + (-1)^3 a_{12} M_{12} + (-1)^4 a_{13} M_{13} \\ &= \underbrace{a_{11}}_2 M_{11} - \underbrace{a_{12}}_{-1} M_{12} + \underbrace{a_{13}}_4 M_{13} \end{aligned}$$

The minor of $a_{11} = 2$ is:

$$M_{11} = \begin{vmatrix} \cancel{2} & \cancel{1} & \cancel{4} \\ \cancel{4} & \mathbf{0} & \mathbf{5} \\ \cancel{-6} & \mathbf{2} & \mathbf{3} \end{vmatrix} = \begin{vmatrix} 0 & 5 \\ 2 & 3 \end{vmatrix} = (0 \times 3) - (5 \times 2) = -10$$

The minor of $a_{12} = -1$ is:

$$M_{12} = \begin{vmatrix} \cancel{2} & \cancel{1} & \cancel{4} \\ \mathbf{4} & \emptyset & \mathbf{5} \\ -6 & \cancel{2} & \mathbf{3} \end{vmatrix} = \begin{vmatrix} 4 & 5 \\ -6 & 3 \end{vmatrix} = (4 \times 3) - (5 \times -6) = 42$$

The minor of $a_{13} = -1$ is:

$$M_{13} = \begin{vmatrix} 4 & 0 \\ -6 & 2 \end{vmatrix} = (4 \times 2) - (0 \times -6) = 8$$

All together:

$$\begin{aligned} |A| &= 2 \times (-10) - (-1) \times 42 + 4 \times 8 \\ &= -20 + 42 + 32 \\ &= 54 \end{aligned}$$

Linear combination

The first two rows are a linear combination, row 1 multiplied by 2 is row 2. Therefore the determinant is zero!

$$\begin{vmatrix} 1 & 2 & 1 \\ 2 & 4 & 2 \\ 3 & -1 & 0 \end{vmatrix} = 0$$

All elements in the lower/upper triangle are zero

The determinant is the product of the leading diagonal.

$$\begin{vmatrix} \mathbf{1} & 0 & 0 & 0 \\ 2 & \mathbf{4} & 0 & 0 \\ 5 & 8 & \mathbf{9} & 0 \\ 20 & -4 & 3 & \mathbf{-1} \end{vmatrix} = 1 \times 4 \times 9 \times (-1) = -36$$

$$\begin{vmatrix} 1 & 4 & 2 & 8 \\ 0 & 4 & 5 & 6 \\ 0 & 0 & 9 & 1 \\ 0 & 0 & 0 & -1 \end{vmatrix} = 1 \times 4 \times 9 \times (-1) = -36$$

1.6 Adjoint

1.6.1 Definition and properties

The adjoint of a square matrix A is the **transpose of the matrix of cofactors**, so for a 3×3 matrix A :

$$\text{adj}(A) = \begin{pmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \\ A_{31} & A_{32} & A_{33} \end{pmatrix}^T$$

Reminder: Cofactor $A_{ij} = (-1)^{i+j} M_{ij}$ (appropriate sign \times minor).

- $A(\text{adj } A) = |A|I$
- $\text{adj}(AB) = (\text{adj } B)(\text{adj } A)$

1.6.2 Example(s)

$$A = \begin{pmatrix} 1 & -2 & 0 \\ 3 & 1 & 5 \\ -1 & 2 & 3 \end{pmatrix} = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$$

Cofactors for the first row:

$$\begin{aligned}A_{11} &= (-1)^{1+1}M_{11} = M_{11} \\&= \begin{vmatrix} 1 & 5 \\ 2 & 3 \end{vmatrix} = 1 \times 3 - 5 \times 2 \\&= -7 \\A_{12} &= (-1)^{1+2}M_{12} = -M_{12} \\&= - \begin{vmatrix} 3 & 5 \\ -1 & 3 \end{vmatrix} = -(3 \times 3 - 5 \times (-1)) \\&= -14 \\A_{13} &= (-1)^{1+3}M_{13} = M_{13} \\&= \begin{vmatrix} 3 & 1 \\ -1 & 2 \end{vmatrix} = 3 \times 2 - 1 \times (-1) \\&= 7\end{aligned}$$

Cofactors for the second row:

$$\begin{aligned}A_{21} &= (-1)^{2+1}M_{21} = -M_{21} \\&= - \begin{vmatrix} -2 & 0 \\ 2 & 3 \end{vmatrix} = -(-2 \times 3 - 0 \times 2) \\&= 6 \\A_{22} &= (-1)^{2+2}M_{22} = M_{22} \\&= \begin{vmatrix} 1 & 0 \\ -1 & 3 \end{vmatrix} = 1 \times 3 - 0 \times -1 \\&= 3 \\A_{23} &= (-1)^{2+3}M_{23} = -M_{23} \\&= - \begin{vmatrix} 1 & -2 \\ -1 & 2 \end{vmatrix} = -(1 \times 2 - (-2) \times (-1)) \\&= 0\end{aligned}$$

Cofactors for the third row:

$$\begin{aligned}
 A_{31} &= (-1)^{3+1}M_{31} = M_{31} \\
 &= \begin{vmatrix} -2 & 0 \\ 1 & 5 \end{vmatrix} = (-2) \times 5 - 0 \times 1 \\
 &= -10
 \end{aligned}$$

$$\begin{aligned}
 A_{32} &= (-1)^{3+2}M_{32} = -M_{32} \\
 &= -\begin{vmatrix} 1 & 0 \\ 3 & 5 \end{vmatrix} = -(1 \times 5 - 0 \times 3) \\
 &= -5
 \end{aligned}$$

$$\begin{aligned}
 A_{33} &= (-1)^{3+3}M_{33} = M_{33} \\
 &= \begin{vmatrix} 1 & -2 \\ 3 & 1 \end{vmatrix} = 1 \times 1 - (-2) \times 3 \\
 &= 7
 \end{aligned}$$

$$\text{adj}(A) = \begin{pmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \\ A_{31} & A_{32} & A_{33} \end{pmatrix}^T = \begin{pmatrix} -7 & -14 & 7 \\ 6 & 3 & 0 \\ -10 & -5 & 7 \end{pmatrix}^T = \begin{pmatrix} -7 & 6 & -10 \\ -14 & 3 & -5 \\ 7 & 0 & 7 \end{pmatrix}$$

1.7 Inverse matrices

1.7.1 Definition and properties

The inverse of a matrix A is given by:

$$A^{-1} = \frac{\text{adj}(A)}{\det(A)} = \frac{\text{adj}(A)}{|A|}$$

- If A is **non-singular** (i. e. $|A| \neq 0$) the inverse of A exists.
- If A is **singular** (i. e. $|A| = 0$) the inverse of A **does not** exist.
- $(AB)^{-1} = B^{-1}A^{-1}$

1.8 Eigenvalue

1.8.1 Definition and properties

Given a square matrix A , its eigenvalues are the **roots** of the following polynomial (**characteristic equation**). These are the values of λ for which $AX = \lambda X$ has no-trivial solutions.

$$p(\lambda) = \det(\lambda I - A)$$

- The denominator of the **transfer function** of a system is equal to the characteristic equation $p(\lambda) = \det(\lambda I - A)$. Therefore the poles of a transfer function are equal to the eigenvalues of the matrix A .

1.8.2 Example(s)

$$A = \begin{pmatrix} 1 & 2 \\ -1 & 4 \end{pmatrix}$$

$$\begin{aligned} p(\lambda) &= |\lambda I - A| = \left| \lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - \begin{pmatrix} 1 & 2 \\ -1 & 4 \end{pmatrix} \right| \\ &= \left| \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix} - \begin{pmatrix} 1 & 2 \\ -1 & 4 \end{pmatrix} \right| = \left| \begin{pmatrix} \lambda - 1 & -2 \\ 1 & \lambda - 4 \end{pmatrix} \right| \\ &= (\lambda - 1)(\lambda - 4) + 2 = \lambda^2 - 4\lambda - \lambda + 4 + 2 \\ &= \lambda^2 - 5\lambda + 6 \end{aligned}$$

Roots:

$$\begin{aligned} p(\lambda) &= \lambda^2 - 5\lambda + 6 = 0 \\ &\rightarrow \lambda_1 = 3 \\ &\rightarrow \lambda_2 = 2 \\ &\rightarrow p(\lambda) = (\lambda - 3)(\lambda - 2) \end{aligned}$$

Eigenvalues of A are 3 and 2.

2 Linear equations

A system of simultaneous linear equations can be written in matrix notation.

$$\begin{aligned} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n &= b_1 \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n &= b_2 \\ \vdots &= \vdots \\ a_{n1}x_1 + a_{n2}x_2 + \dots + a_{nn}x_n &= b_n \end{aligned}$$

Or:

$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{2n} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix}$$

Or:

$$Ax = b$$

- If $b = 0$ the equations are called **homogeneous**.
- If $b \neq 0$ the equations are called **non-homogeneous/inhomogeneous**.
- Cases:
 1. $b \neq 0$ and $|A| \neq 0$: One solution $x = A^{-1}b$.
 2. $b = 0$ and $|A| \neq 0$: Trivial solution $x = 0$.
 3. $b \neq 0$ and $|A| = 0$: No solution **or** infinitely many solutions.
 4. $b = 0$ and $|A| = 0$: Infinitely many solutions.

3 Laplace transformation

3.1 Definition

$$G(s) = \int_0^{\infty} g(t) \cdot e^{-st} dt = \mathcal{L}\{g(t)\} \quad (1)$$

With s as a complex variable and $g(t) = 0$ for $t < 0$!

3.2 A few important properties

If

$$F(s) = \mathcal{L}\{f(t)\}$$

then

$$\mathcal{L}\left\{\frac{df(t)}{dt}\right\} = sF(s) - f(0)$$

(with $f(0)$ =initial condition) and

$$\mathcal{L}\left\{\int f(t)dt\right\}.$$

Finally,

$$\mathcal{L}\left\{\frac{d^n f(t)}{dt^n}\right\} = s^n F(s)$$

if the initial conditions are zero. When we calculate the transfer function of a system we always assume the initial conditions to be zero!

3.3 Table of important transforms

With $n \in \mathbb{N}$ and $a \in \mathbb{C}$. Furthermore, if not noted differently, $\operatorname{Re}(s) > 0$.

$g(t)$ with $t > 0$	$G(s)$
1 (step function)	$\frac{1}{s}$
$\delta(t)$ (Dirac function)	1
e^{at}	$\frac{1}{s-a}$ für $\operatorname{Re}(s) > \operatorname{Re}(a)$
t^n	$\frac{n!}{s^{(n+1)}}$
$t^n \cdot e^{at}$	$\frac{n!}{(s-a)^{n+1}}$ für $\operatorname{Re}(s) > \operatorname{Re}(a)$
$\sin(\omega t)$	$\frac{\omega}{s^2 + \omega^2}$
$\cos(\omega t)$	$\frac{s}{s^2 + \omega^2}$
$t \cdot \sin(\omega t)$	$\frac{2\omega s}{(s^2 + \omega^2)^2}$
$t \cdot \cos(\omega t)$	$\frac{s^2 - \omega^2}{(s^2 + \omega^2)^2}$
$\frac{k\omega_0^2}{s^2 + 2D\omega_0 \cdot s + \omega_0^2}$	$\frac{k \cdot \omega_0 \cdot e^{-D\omega_0 t}}{\sqrt{1-D^2}} \cdot \sin(\omega_0 t \sqrt{1-D^2})$ $0 \leq D < 1$
$\frac{k\omega_0^2}{s(s^2 + 2D\omega_0 \cdot s + \omega_0^2)}$	$k \left[1 - e^{-D\omega_0 t} \left(\cos(\omega_0 t \sqrt{1-D^2}) + \frac{D}{\sqrt{1-D^2}} \sin(\omega_0 t \sqrt{1-D^2}) \right) \right]$ $0 \leq D < 1$

4 State space

4.1 How to transform a n th order differential equation into a set of first order differential equations

Consider the n th order differential equation (input-output relationship/description):

$$a_n \frac{d^n y(t)}{dt^n} + a_{n-1} \frac{d^{n-1} y(t)}{dt^{n-1}} + \dots + a_2 \frac{d^2 y(t)}{dt^2} + a_1 \frac{dy(t)}{dt} + a_0 y(t) = b_0 u(t) \quad (2)$$

With: $a_n, a_{n-1}, \dots, a_0 = \text{constant}$ (parameters)

→ LTI-System (Linear Time Invariant)

Since the system is of order n , we need n states variables: x_1, x_2, \dots, x_n

The question is: How to we define x_1, x_2, \dots, x_n ? We set:

$$\begin{aligned} x_1(t) &= y(t) \\ x_2(t) &= \frac{d^2 y(t)}{dt^2} = \dot{y}(t) = \dot{x}_1(t) \\ x_3(t) &= \frac{d^3 y(t)}{dt^3} = \ddot{y}(t) = \dot{x}_2(t) \\ x_4(t) &= \frac{d^4 y(t)}{dt^4} = \ddot{\dot{y}}(t) = \dot{x}_3(t) \\ x_5(t) &= \frac{d^5 y(t)}{dt^5} = y^{(4)}(t) = \dot{x}_4(t) \\ &\vdots \\ x_{n-1}(t) &= \frac{d^{n-2} y(t)}{dt^{n-2}} = y^{(n-2)}(t) = \dot{x}_{(n-2)}(t) \\ x_n(t) &= \frac{d^{n-1} y(t)}{dt^{n-1}} = y^{(n-1)}(t) = \dot{x}_{(n-1)}(t) \end{aligned}$$

So far we have:

$$\begin{aligned} y(t) &= x_1(t) \\ x(t) &= \begin{pmatrix} x_1(t) \\ x_1(t) \\ \vdots \\ x_n(t) \end{pmatrix} \end{aligned}$$

What we need is: $\dot{x}(t)$. Let's have a look on (2), for the sake of simplicity/clearness (t) is left out:

$$a_n \underbrace{y^{(n)}}_{\dot{x}_n} + a_{n-1} \underbrace{y^{(n-1)}}_{x_n = \dot{x}_{n-1}} + \dots + a_2 \underbrace{\ddot{y}}_{x_3 = \dot{x}_2} + a_1 \underbrace{\dot{y}}_{x_2 = \dot{x}_1} + a_0 \underbrace{y^{(0)}}_{x_1} = b_0 u$$

Now we can write for the highest derivation $\rightarrow y^{(n)} = \dot{x}_n$ (which is not a state):

$$y^{(n)}(t) = -\frac{a_0}{a_n} y(t) - \frac{a_1}{a_n} \dot{y}(t) - \frac{a_2}{a_n} \ddot{y}(t) - \dots - \frac{a_{n-1}}{a_n} y^{(n-1)}(t) + \frac{b_0}{a_n} u(t) \quad (3)$$

Or:

$$\dot{x}_n(t) = -\frac{a_0}{a_n} x_1(t) - \frac{a_1}{a_n} x_2(t) - \frac{a_2}{a_n} x_3(t) - \dots - \frac{a_{n-1}}{a_n} x_n(t) + \frac{b_0}{a_n} u(t) \quad (4)$$

In Summary we have:

$$\begin{aligned} \dot{x}_1(t) &= x_2(t) \\ \dot{x}_2(t) &= x_3(t) \\ \dot{x}_3(t) &= x_4(t) \\ &\vdots \\ \dot{x}_{n-1}(t) &= x_n(t) \\ \dot{x}_n(t) &= -\frac{a_0}{a_n} x_1(t) - \frac{a_1}{a_n} x_2(t) - \frac{a_2}{a_n} x_3(t) - \dots - \frac{a_{n-1}}{a_n} x_n(t) + \frac{b_0}{a_n} u(t) \\ y(t) &= x_1(t) \end{aligned}$$

Or in state space form, again, for the sake of simplicity/clearness (t) is left out:

$$\begin{aligned} \underbrace{\begin{pmatrix} \dot{x}_1 \\ \dot{x}_2 \\ \dot{x}_3 \\ \vdots \\ \dot{x}_{n-1} \\ \dot{x}_n \end{pmatrix}}_{\dot{x}} &= \underbrace{\begin{pmatrix} 0 & 1 & 0 & 0 & \dots & 0 \\ 0 & 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & 0 & 0 & 0 & 1 \\ -\frac{a_0}{a_n} & -\frac{a_1}{a_n} & -\frac{a_2}{a_n} & \dots & -\frac{a_{n-2}}{a_n} & -\frac{a_{n-1}}{a_n} \end{pmatrix}}_{A \text{ (} A \text{ is of the } \textit{companion form} \text{ (german: Kardinalform).)} } \underbrace{\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_{n-1} \\ x_n \end{pmatrix}}_x + \underbrace{\begin{pmatrix} 0 \\ 0 \\ 0 \\ \vdots \\ 0 \\ \frac{b_0}{a_n} \end{pmatrix}}_B u \\ y &= \underbrace{\begin{pmatrix} 1 & 0 & \dots & 0 \end{pmatrix}}_C \underbrace{\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_{n-1} \\ x_n \end{pmatrix}}_x \end{aligned}$$

4.2 Solution of a linear system

$$\dot{x}(t) = Ax(t) + Bu(t)$$

$$y(t) = Cx(t)$$

$x \in \mathbb{R}^n; u \in \mathbb{R}$ valid mostly for single input systems

First of all note:

$$X(s) = (sI - A)^{-1}x(0) + \underbrace{(sI - A)^{-1}BU(s)}_{\text{convolution}}$$

By taking the Laplace inverse of the above equation, we get:

$$x(t) = e^{At}x(0) + \int_0^t e^{A(t-\tau)}Bu(\tau)d\tau \quad (5)$$

With:

$$e^{At} = \mathcal{L}^{-1} \{ (sI - A)^{-1} \}$$

In more general term (for general initial conditions),

$$x(t) = e^{A(t-t_0)}x(t_0) + \int_{t_0}^t e^{A(t-\tau)}Bu(\tau)d\tau \quad (6)$$

4.3 Transformation from state-space to transfer function

$$\sum \begin{cases} \dot{x}(t) & = Ax(t) + Bu(t) \\ y(t) & = Cx(t) \end{cases}$$

$x \in \mathbb{R}^n; u \in \mathbb{R}$ valid mostly for single input systems

By taking the Laplace transform of system (\sum) we get:

$$\mathcal{L} \{ \dot{x}(t) \} = sX(s) - x(0)$$

With:

$$X(s) = \mathcal{L} \{ x(t) \} = \begin{pmatrix} \mathcal{L} \{ x_1(t) \} \\ \mathcal{L} \{ x_2(t) \} \\ \vdots \end{pmatrix} = \begin{pmatrix} X_1(s) \\ X_2(s) \\ \vdots \end{pmatrix}$$

Hence:

$$sX(s) - x(0) = AX(s) + BU(s) \quad (7)$$

$$Y(s) = CX(s) \quad (8)$$

From (7), we have:

$$\begin{aligned} sX(s) - AX(s) &= x(0) + BU(s) \\ (sI - A)X(s) &= x(0) + BU(s) \\ X(s) &= (sI - A)^{-1}(x(0) + BU(s)) \end{aligned}$$

From (8), we have:

$$Y(s) = CX(s) = C(sI - A)^{-1}x(0) + C(sI - A)^{-1}BU(s) \quad (9)$$

When we calculate a transfer function we always assume $x(0) = 0$. Hence,

$$Y(s) = C(sI - A)^{-1}BU(s)$$

That is:

$$H(s) = \frac{Y(s)}{U(s)} = \frac{\text{output}}{\text{input}} = C(sI - A)^{-1}B \quad (10)$$

Reminder: $U(s)$ is not a matrix because $u \in \mathbb{R}$ (single input system).

5 From the input-output relationship to the transfer function

Let

$$Y(s) = \mathcal{L}\{y(t)\}$$

and

$$U(s) = \mathcal{L}\{u(t)\}.$$

Then taking the Laplace transform of the input-output relationship (2) given above, we have:

$$a_n s^n Y(s) + a_{n-1} s^{n-1} Y(s) + \dots + a_2 s^2 Y(s) + a_1 s Y(s) + a_0 Y(s) = b_0 U(s)$$

Or:

$$Y(s)[a_n s^n + \dots + a_0] = b_0 U(s)$$

Therefore:

$$H(s) = \frac{Y(s)}{U(s)} = \frac{b_0}{a_n s^n + a_{n-1} s^{n-1} + \dots + a_2 s^2 + a_1 s + a_0}$$